

Nuclear stellar discs in low-luminosity elliptical galaxies: NGC 4458 and NGC 4478

L. Morelli^{1,2}, C. Halliday³, E. M. Corsini¹, A. Pizzella¹, D. Thomas⁴, R. P. Saglia⁴,
 R. L. Davies⁵, R. Bender^{4,6}, M. Birkinshaw⁷, and F. Bertola¹

¹ Dipartimento di Astronomia, Università di Padova, vicolo dell'Osservatorio 2, I-35122 Padova, Italy.

² European Southern Observatory, 3107 Alonso de Cordova, Santiago, Chile.

³ INAF-Osservatorio Astronomico di Padova, vicolo dell'Osservatorio 5, I-35122 Padova, Italy.

⁴ Max-Planck Institut für extraterrestrische Physik, Giessenbachstrasse, D-85748 Garching, Germany.

⁵ Department of Astrophysics, University of Oxford, Keble Road, Oxford, OX1 3RH.

⁶ Universitäts-Sternwarte, Scheinerstrasse 1, D-81679 Muenchen, Germany.

⁷ H. H. Wills Physics Laboratory, University of Bristol, Tyndall Avenue, Bristol BS8 1TL, UK.

Received 2004 May 20; accepted 2004 July 21

ABSTRACT

We present the detection of nuclear stellar discs in the low-luminosity elliptical galaxies NGC 4458 and NGC 4478, which are known to host a kinematically-decoupled core. Using archival HST imaging, and available absorption line-strength index data based on ground-based spectroscopy, we investigate the photometric parameters and the properties of the stellar populations of these central structures. Their scale length, h , and face-on central surface brightness, μ_0^c , fit on μ_0^c-h relation for galaxy discs. For NGC 4458 these parameters are typical for nuclear discs, while the same quantities for NGC 4478 lie between those of nuclear discs and the discs of discy ellipticals. We present Lick/IDS absorption line-strength measurements of $H\beta$, $Mg\ b$, $\langle Fe \rangle$ along the major and minor axes of the galaxies. We model these data with simple stellar populations that account for the α/Fe overabundance. The counter-rotating central disc of NGC 4458 is found to have similar properties to the decoupled cores of bright ellipticals. This galaxy has been found to be uniformly old despite being counter-rotating. In contrast, the cold central disc of NGC 4478 is younger, richer in metals and less overabundant than the main body of the galaxy. This points to a prolonged star formation history, typical of an undisturbed disc-like, gas-rich (possibly pre-enriched) structure.

Key words: galaxies: elliptical and lenticular, cD — galaxies: photometry — galaxies: kinematics and dynamics — galaxies: abundances — galaxies: formation — galaxies: evolution

1 INTRODUCTION

In recent years, the sub-arcsec resolution of the *Hubble Space Telescope* has allowed the study of galactic nuclei, unveiling the presence of distinct components such as small-scale stellar discs (see Pizzella et al. 2002 and references therein) and nuclear clusters (Carollo et al. 1997; Böker et al. 2002; Graham & Guzman 2003). To date surface-brightness distributions of nuclear stellar discs (hereinafter NSD) have been measured for only a few of S0s (Scorza & van den Bosch 1998; van den Bosch et al. 1998; Kormendy et al. 1996) and early-type spiral galaxies (Pizzella et al. 2002). They have smaller scale lengths ($\sim 10-30$ pc) and higher central surface brightnesses ($\sim 15-19$ mag arcsec⁻² in the V band) with respect to those of embedded stellar discs, which have been found in several elliptical and lenticular galaxies (e.g., Scorza & Bender 1995, hereinafter SB95). Although only few NSDs have been studied in detail, they may be quite a common structure in

the central regions of spheroids. Indeed, nuclear discs of gas, dust and stars have been clearly detected in a large number of early-type galaxies (e.g., Jaffe et al. 1994; Lauer et al. 1995; van Dokkum & Franx 1995; Faber et al. 1997; Carollo et al. 1997; Tomita et al. 2000; Kormendy et al. 2001; Tran et al. 2001; Trujillo et al. 2004), but the photometric parameters of such discs have not yet been derived. Furthermore, HST imaging in near-infrared band-passes provided indirect signatures of the presence of NSDs as discy isophotes in the nuclei of early-type galaxies (Ravindranath et al. 2001) or photometrically-distinct exponential components in bulges (Balcells et al. 2003, 2004). The existence of NSDs suggests that the continuity of the disc properties, with a smooth variation of scale parameters from spirals to discy ellipticals along a sequence of decreasing disc-to-bulge ratio (Kormendy & Bender 1996), could be extended to nuclear scales (van den Bosch 1998). In this framework NSDs could provide important clues to the assembly scenario of their host galaxies.

In the current picture, the NSDs observed in bulges of S0s and spiral galaxies are believed to have formed either from a secular evolution of a nuclear bar (*e.g.*, NGC 4570, van den Bosch et al. 1998; Scorza & van den Bosch 1998; van den Bosch & Emsellem 1998) or as the end result of a merging event (*e.g.*, NGC 4698, Bertola et al. 1999; Pizzella et al. 2002). Each of these scenarios is likely to be correct for some but not for all the objects. In both of them the gas is efficiently directed toward the galaxy centre, where it first dissipates and settles onto an equilibrium plane and then forms into stars. Although the decoupled kinematics strongly suggest a later infall (*e.g.*, NGC 4486A, Kormendy et al. 2001) or merger (*e.g.*, Holley-Bockelmann & Richstone 2000), the frequently indistinguishable star formation history remains an enigma. Yet, these processes are expected to lead to distinct features in stellar populations of NSDs and surrounding spheroids. The measurement of the kinematics of some early-type galaxies (elliptical and S0 galaxies) revealed evidence for kinematically decoupled or peculiar galaxy cores (see Bertola & Corsini 1999 for a review), strongly indicative of formation by galaxy-galaxy merging (*e.g.*, Kormendy 1984; see Mehlert et al. 1998) and references therein). Such kinematically-distinct cores have been associated with a central disc component (*e.g.*, Rix & White 1992; Mehlert et al. 1998). However, triaxial galaxies with radius-dependent ellipsoids, can, in projection, mimic kinematically decoupled cores (see Arnold, de Zeeuw & Hunter 1994, and references therein). Spectroscopic absorption line-strength indices (Faber et al. 1985; Trager et al. 1998) constrain the luminosity-weighted stellar population age, metallicity and alpha-element overabundance by comparison with evolutionary population synthesis models (Worthey 1994; Maraston 1998; Tantalo et al 1998; Vazdekis 1999; Thomas, Maraston & Bender 2003, hereafter TMB; Bruzual & Charlot 2003). Absorption line-strength indices have been measured out to one effective radii for elliptical galaxies (*e.g.*, Davies et al. 1993; González 1993; Halliday 1999; Mehlert et al. 2000). Such measurements can distinguish between merging and dissipational formation scenarios by the constraint of metallicity gradients, dating of the latest episode of star formation and the measurement of the timescale over which the bulk of the star formation has taken place. Davies et al. (1993), Mehlert et al. (2003) and the analysis of literature data by Kobayashi & Arimoto (1999) have found elliptical galaxies to have gradients in metallicity shallower than predicted by dissipational collapse models. Mehlert et al. (1998) presented spectroscopic line-strength data and HST archive imaging for two Coma cluster galaxies with kinematically-decoupled cores and found little evidence that the cores have experienced different star formation histories from their parent galaxies. Using wide-field spectroscopic maps of NGC 4365 acquired using the SAURON instrument, Davies et al. (2001) presented evidence for a metal-rich core but constant age and α /Fe-enhancement for both the kinematically-distinct core and parent galaxy. Both studies are consistent with the core and the main galaxy having experienced similar star formation histories.

In this paper, we use archival HST imaging and ground-based spectroscopy to investigate the photometric parameters and stellar population diagnostics of the NSDs hosted by the two Virgo galaxies NGC 4458 and NGC 4478. NGC 4458 and NGC 4478 are classified as E0-1 (de Vaucouleurs et al. 1991, hereafter RC3) and E2 (Sandage & Tammann 1981; RC3), respectively. They are both low-luminosity ellipticals with a $M_B^0 \sim -18$ (RC3) at a distance of 12.6 Mpc (Tully 1988, $H_0 = 100 \text{ km s}^{-1} \text{ Mpc}^{-1}$). They have a power-law central luminosity profile (Ferrarese et al. 1994; Faber et al. 1997; Rest et al. 2001) and both galaxies show deviations

from the $r^{1/4}$ law in their outskirts (Michard 1985; Prugniel et al. 1987; Caon et al. 1990; Peletier et al. 1990), which have been explained as due to tidal interaction with NGC 4461 and NGC 4486. Recently, the stellar kinematics along the major and minor axes of NGC 4458 and NGC 4478 have been measured by Halliday et al. (2001), who found the signature of a kinematically-decoupled core in their inner ~ 5 arcsec. In particular, NGC 4458 has a clear counter-rotating core along the major axis, while NGC 4478 has a cold component detected along both the major and the minor axis.

In Section 2 we perform the photometric analysis of the archival HST images of the nuclei of NGC 4458 and NGC 4478 in order to derive the surface brightness profiles and scale parameters of their NSDs. In Section 3 we present the absorption line-strength indices as a function of radius of the major and minor axes of both galaxies from the thesis research of Halliday (1999). In Section 4 we combine photometric and spectroscopic results to assess age, metallicity and overabundances of stellar populations of the two NSDs and surrounding spheroids. This allowed us to suggest possible formation and evolution scenarios for NGC 4458 and NGC 4478. We summarize our conclusions in Section 5.

2 PHOTOMETRIC PARAMETERS

2.1 Data reduction

We retrieved Wide Field Planetary Camera 2 (WFPC2) images of NGC 4458 and NGC 4478 from the HST archive. Data for the filter F814W were selected as a compromise between obtaining data in identical bandpasses for both galaxies and minimizing the effects of dust on photometric measurements. Total exposure times were 1120 s for NGC 4458 (5 exposures, Prog. Id. 5512, P.I. S. M. Faber) and 1600 s for NGC 4478 (2 exposures, Prog. Id. 6587, P.I. D. Richstone), respectively. All exposures were taken with the telescope guiding in fine lock, which typically gave an rms tracking error of 0.003 arcsec. We focused our attention on the Planetary Camera chip (PC) where the nucleus of both galaxies was centred. This consists of 800×800 pixels of $0.0455 \times 0.0455 \text{ arcsec}^2$ each, yielding a field of view of about $36 \times 36 \text{ arcsec}^2$. The images were calibrated using the standard reduction pipeline maintained by the Space Telescope Science Institute. Reduction steps include bias subtraction, dark current subtraction, and flat-fielding and are described in detail in Holtzman et al. (1995a). Subsequent reduction was completed using standard tasks in the STSDAS package of IRAF¹. Bad pixels were corrected by means of a linear one-dimensional interpolation using the data quality files and the WFIXUP task. Different images of the same target were aligned and combined using IMSHIFT and knowledge of the offset shifts. Cosmic ray events were removed using the task CRREJ. The cosmic-ray removal and bad pixel correction were checked by inspection of the residual images between the cleaned and combined image and each of the original frames. Residual cosmic rays and bad pixels in the PC were corrected by manually editing the combined image with IMEDIT. The sky level ($\sim 1 \text{ count pixel}^{-1}$) was determined from regions free of sources in the Wide Field chips and subtracted from the PC frame after appropriate scaling.

¹ IRAF is distributed by NOAO, which is operated by AURA Inc., under contract with the National Science Foundation




Figure 1. *Left panels:* WFPC2/F814W images of NGC 4458 and NGC 4478 (for which nuclear discs were detected), and NGC 4539 (for which a nuclear disc was not detected). The size of the plotted region is 19.3×19.3 arcsec². The orientation is specified by the arrow indicating north and the segment indicating east in the lower right corner of each panel. *Middle and right panels:* Unsharp masking of the WFPC2/F814W images obtained with $\sigma = 2$ and 6 pixels, respectively. Sizes and orientations are as in the left-hand panels.

2.2 Detection of nuclear discs

To test for the presence of a NSD in NGC 4458 and NGC 4478, we constructed the unsharp-masked image of the PC frame using an identical procedure to Pizzella et al. (2002). We divided each image by itself after convolution by a circular Gaussian of width $\sigma = 2$ and 6 pixels, corresponding to 0.09 arcsec and 0.27 arcsec, respectively (Figure 1). This procedure enhanced any surface-brightness fluctuation and non-circular structure extending over a spatial region comparable to the σ of the smoothing Gaussian. Two values of σ were adopted to attempt the identification of structures with different scale lengths.

A highly elongated structure is clearly visible in the nucleus of each galaxies. These structures are not artifacts of the unsharp-masking procedure. In Figure 1 we show as a counter-example the unsharp-masked image of NGC 4539 which has been obtained using an identical method to that for NGC 4458 and NGC 4478; we clearly do not detect an elongated disc-like structure. Moreover the nuclear structures of NGC 4458 and NGC 4478 are associated with a central increase in ellipticity measured by performing an isophotal analysis using the IRAF task ELLIPSE (Figure 2).

We analyzed the isophotal profiles of both galaxies by first masking foreground stars and then fitting ellipses to the isophotes. We allowed the centres of the ellipses to vary, to test whether the galaxies were disturbed. Within the errors of the fits, we found no evidence of variations in the fitted centre. The ellipse fitting was repeated with the ellipse centres fixed. The resulting azimuthally averaged surface brightness, ellipticity, position angle, the fourth (a_4), and sixth (a_6) cosine Fourier coefficients profiles are presented in Figure 2. In the innermost ~ 2 arcsec we measured positive values of the a_4 , and a_6 Fourier coefficients, which describe




Figure 2. (a) Radial profiles of surface brightness of NGC 4458 (left panel) and NGC 4478 (right panel) after deconvolution (open squares) are compared to those of their nuclear discs (thick lines). (b) Radial profiles of the fraction of total luminosity contributed by the nuclear disc of NGC 4458 (left panel) and NGC 4478 (right panel). Radial profiles of (c) ellipticity, (d) position angle, (e) a_4 and (f) a_6 Fourier coefficients before (open squares) and after (filled circles) the subtraction of the best-fitting model for the nuclear disc of NGC 4458 (left panels) and NGC 4478 (right panels).

the discy deviation of the isophotes from pure ellipses (Jedrzejewski 1987). These photometric features have been also observed in the HST/F555W image for NGC 4458 (Lauer et al. 1995; Trujillo et al. 2004) and in the HST/F702W image of NGC 4478 (van den Bosch et al. 1994; Rest et al. 2001; Trujillo et al. 2004) and confirm the presence of a NSD in both galaxies.

2.3 Photometric decomposition

2.3.1 Photometric parameters of the nuclear discs with the Scorza & Bender method

After establishing the existence of NSDs, we measured their photometric properties using the method described by SB95. This method is based on the assumption that isophotal disciness is the result of the superimposition of a spheroidal component (which is either a host elliptical galaxy or a bulge component) and an inclined disc. The two components are assumed to have both perfectly elliptical isophotes with constant but different ellipticities. When adopting this technique to study the innermost regions of galaxies it is essential to restore the images from the effects of the HST point spread function (PSF) in order to properly derive the photometric parameters of the NSDs, as shown by Scorza &

van den Bosch (1998). Such deconvolution was performed through the Richardson-Lucy method by means of the IRAF task LUCY. Although susceptible to noise amplification, this algorithm was shown by van den Bosch et al. (1998) to lead to a restored surface-brightness distribution comparable to that obtained using a multi-Gaussian representation (Monnet et al. 1992; Cappellari 2002). We believe that the results obtained by van den Bosch et al. (1998) are directly applicable to our case: we are dealing with images obtained with similar or longer integration times, galaxies with less steep surface-brightness profiles and NSDs with equal or larger scale lengths. We decided to deconvolve the images with a number of iterations between 3 and 6. A larger number of iterations does not affect the result of the decomposition but does amplify the noise. For each given image and nucleus position on the PC we adopted a model PSF calculated using the TINY-TIM package (Krist & Hook 1999). No correction for telescope jitter was necessary. The SB95 method consists of the iterative subtraction of a thin disc model characterized by an exponential surface-brightness profile with central surface brightness I_0 and radial scale length h , and by an axial ratio b/a . The disc parameters are adjusted until the departures from perfect ellipses are minimized (*i.e.*, a_4 and a_6 are close to zero). For each disc model we obtained the disc-free image of the galaxy by subtracting the disc model from the galaxy image. We performed an isophotal analysis on the disc-free image using the IRAF task ELLIPSE. We defined

$$\chi^2 = \sum_{i=1}^N \frac{a_{4,disc-free}^2(i)}{\sigma(i)^2} \quad (1)$$

where $a_{4,disc-free}(i) \pm \sigma(i)$ is the value of the a_4 Fourier coefficient measured for the i -th isophote in the disc-free image, and N is the number of fitted isophotes. We assumed $\sigma(i) = 0.01$ as typical error on $a_{4,disc-free}$ for all the isophotes in the region of the NSD. We defined the reduced χ^2 as $\chi_v^2 = \chi^2/(N - M)$, where $M = 3$ is the number of free parameters, namely I_0 , h , and b/a . According to our technique for photometric decomposition the minimum value of $\chi_v^2 \equiv \chi_{v,min}^2$ corresponds to the best-fitting model of the NSD. We found $\chi_{v,min}^2 = 0.43$ for NGC 4458 and $\chi_{v,min}^2 = 0.34$ for NGC 4478. We determined $\Delta\chi_v^2 \equiv \chi_v^2 - \chi_{v,min}^2$ and derived its confidence levels under the assumption that the errors are normally distributed (Press et al. 1992). The resulting contour plots of χ_v^2 are shown in Figure 3 and they allowed us to derive the best-fitting values of I_0 , h and b/a and their 1σ errors.

We derived the Johnson V -band central surface brightness of the NSDs from the HST VEGAMAG system (Holtzman et al. 1995b) using the STSDAS task SYNPHOT. Since this correction depends on the galaxy spectral energy distribution, it has been calculated using the spectral template for elliptical galaxies by Kinney et al. (1996). The resulting relation is $V - m_{F814W} = 1.33$. We derived the scale length of the NSDs assuming a distance of 12.6 Mpc for both galaxies. The inclination of NSDs has been calculated as $i = \arccos(b/a)$. The resulting values of central surface brightness in the V band, scale length and inclination for both NSDs are listed in Table 1.

The disciness of NGC 4458 and NGC 4478 disappears after the subtraction of the best-fitting model of the NSD as it results from the elliptical shape of their isophotes as shown by the photometric profiles plotted in Figure 2. We tested that the result of the photometric decomposition is independent of position on the PC chip where the PSF is built. The resulting fraction of total luminosity contributed by the nuclear disc of NGC 4458 and NGC 4478 is plotted as a function of radius in Figure 2.

Table 1. Photometric parameters of the nuclear discs

Name	$\mu_{0,V}$ [mag arcsec ⁻²]	h [pc]	i [°]	L_{disc} [L_{\odot}, V]
NGC 4458	14.74 ^{+0.71} _{-0.67}	11.0 ^{+5.9} _{-2.5}	83.1 ^{+6.9} _{-8.8}	4.8 · 10 ⁶
NGC 4478	16.61 ^{+1.48} _{-0.81}	40.5 ^{+33.0} _{-18.7}	77.9 ^{+11.6} _{-10.0}	1.9 · 10 ⁷

2.3.2 Photometric parameters of the nuclear discs with the Seifert & Scorza method

In applying the decomposition method by SB95 we assumed that the surface-brightness profile of the NSDs was exponential. To verify this assumption, we derived the surface-brightness profile of the NSDs using the alternative method of Seifert & Scorza (1996, hereinafter SS96). This decomposition method was developed to identify embedded discs in ellipticals and bulges without any *a priori* assumptions of the parametric law for their surface-brightness profile. In the deconvolved image of the galaxy we masked the region of the NSD with a double-cone mask, and fit ellipses to the isophotes in the remaining part of image using the IRAF task ELLIPSE to derive the photometric properties of the host galaxy. A double-cone mask centred on the galaxy major axis is an optimal choice for the highly-inclined NSDs of NGC 4458 and NGC 4478 since we expect that the isophotes close to the galaxy minor axis will be only slightly contaminated by light from the NSDs. The very central portion of the galaxy image was not masked to allow meaningful isophotes to be fitted. These masks correspond to the central 3×3 pixel (0.14×0.14 arcsec²) of NGC 4458 and 5×5 pixel (0.23×0.23 arcsec²) of NGC 4478. The opening angle of the mask ($\approx 90^\circ$) was adjusted to minimize the influence of the NSD on the fitted parameters of the surrounding galaxy bulge component, but to ensure that a sufficient number of data points for the model derivation remained. A model was constructed to have perfectly elliptical isophotes with the same surface brightness, ellipticity and position angle radial profiles which we obtained from the isophotal fit. We subtracted the galaxy model from the original image to obtain a residual image which shows an elongated structure, namely the NSD, responsible for the higher-order deviations from perfectly elliptical isophotes. We extracted the surface-brightness radial profile along the major axis of the residual disc which was close to being exponential. Scale lengths are consistent within their errors to those measured using the SB95 method, while the central surface brightnesses are somewhat fainter. This discrepancy in the surface brightnesses derived for using the SB95 and SS96 methods is due to the actual shape of the double-cone mask where we excluded the central region in order to allow meaningful isophotes to be fitted. In this way the observed central surface brightness is assigned to the host galaxy and the central surface brightness of the NSD is therefore underestimated.

2.4 Structural parameters of nuclear discs

It is interesting to compare the scale lengths given in Table 1 ($h \approx 0.2''$ for NGC 4458 and $h \approx 0.7''$ for NGC 4478) with the “break” radii found in the major and minor axis kinematics of the two galaxies (Halliday et al. 2001). Counter-rotation of about (30 km s⁻¹) is observed in the inner 4 arcsec of the major axis of NGC 4458. In the case of NGC 4478 the velocity dispersion drops in the central 2 arcsec, where the rotational velocity along the major axis




Figure 3. Contour plots of χ^2_{ν} for NGC 4458 (*upper panels*) and NGC 4478 (*lower panels*) as a function of the central surface brightness, I_0 , scale length, h , and axial ratio, b/a , of the nuclear disc models. *Solid contours* define the formal 68.3%, 95.4%, and 99.7% confidence levels. Intermediate confidence levels are given with *dashed contours*. *Filled circles* indicate actual model calculations. Bi-cubic interpolation is used to calculate at χ^2_{ν} intermediate points. *Crosses* correspond to the best-fitting photometric parameters of the nuclear disc.

drops by $\approx 20 \text{ km s}^{-1}$ from 50 km s^{-1} at 2 arcsec to 30 km s^{-1} at 3 arcsec from the centre to increase again at larger radii.

Clearly, the spatial resolution of the kinematic data (see Section 3 for details) is smearing the signal of the NSDs. Figure 4 shows the position of the NSDs of NGC 4458 and NGC 4478 in the disc μ_0^c-h diagram (adapted from Pizzella et al. 2002; but see also van den Bosch 1998, and Graham 2001). The face-on central surface brightness μ_0^c have been derived from the observed one μ_0 by applying the following inclination correction

$$\mu_0^c = \mu_0 - 2.5 \log(\cos i) \quad (2)$$

and without taking into account any correction for extinction. The (small) scale length and (high) central surface brightness of the disc of NGC 4458 are typical for NSDs. In contrast, the (relatively large) scale length and (relatively low) central surface brightness of the disc of NGC 4478 are in between the typical values found for discy ellipticals and NSDs. No correlations are found with global properties (*e.g.*, luminosity, scale-length, disc-to-bulge ratio) of the host galaxies.

Finally, on a speculative line of thought without presumption of rigourness, it is interesting to correlate the order of magnitude rotational velocities estimated above (30 km s^{-1} for NGC 4458 and 50 km s^{-1} for NGC 4478) with the estimated magnitudes of the discs. We find that discs following the Tully-Fisher relation (as in Haynes et al. 1999) and having the same rotational velocities, are expected to be ≈ 3 mag brighter than the NSDs of NGC 4458 and NGC 4478. In other words, the NSDs of NGC 4458 and NGC 4478 rotate faster than discs of the same luminosity that follow the Tully-Fisher relation. Taken at face value, this might indicate that they behave like test particles embedded in the dominant gravita-

tional potential generated by the main (bulge) body of the galaxies. However, the situation is complicated by the low spatial resolution of the kinematic data. Using the total masses of the discs derived in Section 4, we estimate that maximum rotational velocities of the order of $\approx 30 - 50 \text{ km s}^{-1}$ are generated by the NSDs alone around 0.6-2 arcsec. Clearly, higher resolution kinematic data and proper modelling (including a decomposition of the line-of-sight velocity distributions and seeing convolution) are needed to pin down the issue.

3 LINE-STRENGTH INDICES

Absorption line-strength indices are taken from a study of the stellar populations and kinematics of low-luminosity elliptical galaxies, in the Virgo cluster and nearby groups (Halliday 1999; Halliday et al. 2001). We provide a brief summary of our spectroscopic observations. Long-slit spectroscopic data were obtained using the Blue Channel Spectrograph at the Multiple Mirror Telescope, Arizona (USA). A Loral $3\text{k} \times 1\text{k}$ CCD chip with a pixel size of $15\mu\text{m} \times 15\mu\text{m}$, a grating of $1200 \text{ slits mm}^{-1}$ and a slit width of 1 arcsec were used. The spectral resolution was $\sim 1.5 \text{ \AA}$ FWHM, wavelength range $4555 - 6045 \text{ \AA}$ and spatial scale $0.6 \text{ arcsec pixel}^{-1}$. Seeing varied between 0.5 and 0.8 arcsec during observations. The reader is referred to Halliday (1999) for details of the spectroscopic data reduction and Halliday et al. (2001) for a description of the kinematical analyses.




Figure 4. Disc μ_0^c - h diagram adapted from Pizzella et al. (2002). *Open circles* refer to high surface-brightness spirals, *triangles* to low surface-brightness spirals, *stars* to S0's, *filled circles* to discy ellipticals. *Large squares* correspond to NSDs in elliptical galaxies (*open symbols*) and in S0's and spiral galaxies (*filled symbols*), respectively.

3.1 Measurement of absorption line-strength indices

Line-strength indices were measured as a function of radius to approximately one photometric effective radius along the major and minor axes of each galaxy. One-dimensional spectra were extracted for different luminosity-weighted radii of each axis and line-strength indices were measured for each spectrum using an algorithm developed by G. Bagglely and revised by H. Kuntschner (Halliday 1999). The Lick/IDS system line-strength index definitions of Trager et al. (1998) were adopted. A fuller description of the algorithm is given below.

The principal input was a spectroscopic CCD frame for which a basic data reduction had been completed. Before extracting one-dimensional spectra the galaxy spectrum was checked to be aligned with the rows of the CCD frame. Using the IRAF tasks `APFIND` and `APTRACE` the position of the galaxy centre was traced along the dispersion axis; this step was performed to an accuracy of $\simeq 0.03$ pixels rms. The pixels were resampled such that the galaxy spectrum was centred on the same row pixel position as a function of dispersion.

One-dimensional spectra were extracted by summing rows of the galaxy frame until a signal-to-noise ratio (hereinafter S/N) of 40 per \AA (for the wavelength range ~ 5100 – 5300 \AA) was attained. The dispersion axes of extracted spectra were rebinned to logarithmic intervals of wavelength. Redshifts were measured using the IRAF task `FXCOR`, by cross-correlating each spectrum with a range of one-dimensional de-redshifted stellar spectra. The mean redshift determined for all templates was used to de-redshift each galaxy spectrum. $H\beta$, Mg b , Fe5270 and Fe5335 line-strength indices were measured for all spectra. Errors for these raw index measurements were calculated based on Poissonian noise, the typical sky level counts subtracted during sky subtraction, and the values of read-out noise and gain of the CCD.

Our line-strength index measurements were transformed to the Lick/IDS system. This procedure involved two steps: (i) correc-

tion to zero galaxy velocity dispersion and (ii) correction to the Lick/IDS spectral resolution (~ 8.6 \AA).

To correct for the effects of galaxy velocity dispersion, σ , different stellar spectra were smoothed by Gaussians corresponding to measurements of σ between 0 km s^{-1} and 300 km s^{-1} in intervals of 20 km s^{-1} . Smoothing was performed using the IRAF task `GAUSS`. Correction factors for different values of σ were calculated for each index by comparing line index measurements for each smoothed spectrum with measurements for the original unsmoothed stellar spectrum. For the atomic indices $H\beta$, Mg b , Fe5270 and Fe5335 correction factors were defined to be the ratio of measurements for the original unbroadened stellar spectrum, I_{orig} , and that for the spectrum broadened to a particular velocity dispersion, I_σ , *i.e.*,

$$C(\sigma) = \frac{I_{orig}}{I_\sigma}. \quad (3)$$

Mg b , Fe5270 and Fe5335 line-strength index measurements were corrected for the effects of σ using the calibrations for 12 stellar observations; for $H\beta$ data 8 stellar observations were used. Linear relations were interpolated between the mean correction factor for each σ . The correction factor for an arbitrary measurement of σ was determined using the linear relation for the σ range bracketing the σ measurement. The measurement of σ as a function of radius was presented for both galaxies in Halliday et al. (2001). We adopt the σ measurements obtained for a S/N $\simeq 60$ per \AA . Measurements of σ with S/N = 30 per \AA were adopted at large radii where uncertainties were larger.

Absorption line-strength index measurements and measurements of σ from Halliday et al. (2001) were measured in general for different galactic radii. Linear interpolations were made between neighboring measurements of σ . A σ measurement for a radius at which line-strength indices were measured, was found by linear interpolation from bracketing data. Depending on the appropriate side of the galaxy, if the radius was either less than or exceeded the radius of the most reliable measurement of σ , the linear interpolation determined for the closest radial interval was used to determine σ . The errors of line-strength index measurements corrected for galaxy velocity dispersion were calculated to be,

$$\delta I_{atomic}(\sigma) = \sqrt{(\delta I_{orig} \cdot C(\sigma))^2 + (I_{orig} \cdot \delta C(\sigma))^2} \quad (4)$$

where δI_{orig} is the error in the raw measurement calculated as outlined above and $\delta C(\sigma)$ is the addition in quadrature of the standard deviations of correction factors for measurements of σ bracketing the σ measurement.

Corrections for the difference between our spectral resolution and the resolution of the Lick/IDS system were applied. A simulated Lick aperture was extracted for the major axis spectrum of each galaxy studied by Halliday (1999) present in the ‘‘pristine’’ IDS sample of Trager (1997). Spectra were resampled such that the galaxy centre was aligned with the same pixel row position as a function of wavelength. For our spectroscopic slit width of 1 arcsec and spatial scale of 0.6 arcsec pixel $^{-1}$ a one-dimensional spectrum corresponding to a Lick/IDS aperture was created by summing the central 9 rows (*i.e.*, the central ~ 5.4 arcsec of each galaxy spectrum). Line-strength indices were measured using the Lick/IDS definitions of Trager et al. (1998). The galaxy velocity dispersion was measured using the IRAF task `FXCOR`. Line-strength indices were corrected for the effects of σ using the calibrations discussed above. Direct comparisons were made with the fully-corrected Lick/IDS 1.4×4.0 arcsec 2 aperture measurements of the ‘‘pristine’’ IDS sample of Trager (1997); we considered

the Lick/IDS $H\beta$ measurements corrected for the effects of σ but uncorrected for $H\beta$ emission. Relation fitting was attempted between the simulated Lick/IDS aperture measurements corrected for σ and the Lick/IDS measurements of Trager (1997), as a function of our σ -corrected, simulated Lick/IDS aperture measurements. After testing various routines, the median value of offset was found to provide a good fit for all line-strength indices (*i.e.*, for $H\beta$, 0.272; Mg_2 , -0.048; Mg_b , 0.051; Fe5270, 0.544; Fe5335, 0.697). The σ -corrected measurements were established on to the Lick/IDS system by subtracting these median values for all index measurements excluding $Mg\ b$ for which no zero correction was required due to the large scatter of offset values around the median.

3.2 Absorption line-strength gradients

Absorption line-strength index measurements for $H\beta$, $Mg\ b$, Mg_2 and $\langle Fe \rangle = (Fe5270 + Fe5335)/2$ are shown as a function of radius for the major and minor axes of NGC 4458 and NGC 4478 in Figure 5. Line-strength measurements are presented as a function of the normalized radius R/R_e^* , where $R_e^* = R_e/\sqrt{1-\epsilon}$ for the major axis and $R_e^* = R_e \cdot \sqrt{1-\epsilon}$ for the minor axis, respectively. R_e is the circularized effective radius defined as $R_e = A_e/2$ with A_e the diameter of the circle enclosing one-half of the B -band total flux (RC3), and ϵ is the mean ellipticity of the galaxy. We assume $R_e = 26.1$ arcsec (RC3, corresponding to 1.6 kpc at the adopted distance) and $\epsilon = 0.00$ (Lauer et al. 1995) for NGC 4458, $R_e = 13.4$ arcsec (RC3, corresponding to 0.8 kpc at the adopted distance) and $\epsilon = 0.17$ (Peletier et al. 1990) for NGC 4478. The different central values measured along major and minor axis for the same index give an estimate of the true measurement errors. Measurements for Mg_2 are shown for completeness and the calibration of this molecular index to the Lick/IDS system is described in Halliday (1999).

For NGC 4478, measurements of $H\beta$, $Mg\ b$, Mg_2 , Fe5270, Fe5335 and $\langle Fe \rangle$ were derived previously for the galaxy centre by González (1993) and Trager et al. (1998), and as a function of radius of the major axis by Peletier (1989) and Gorgas et al. (1990). In Figure 6 we compare all previous measurements with our measurements presented here.

Good agreement within all margins of uncertainty is found between our data and the measurements of Peletier (1989). Measurements and errors for Fe5270 were not tabulated by Peletier (1989) and have been derived using the corresponding measurements of Fe5335 and $\langle Fe \rangle$. Agreement within the errors is found between our measurements and those of Gorgas et al. (1990) for $H\beta$ and $Mg\ b$ but not for Fe5270, Fe5335, and $\langle Fe \rangle$. A single bad pixel column was interpolated for a region corresponding to the red passband of Fe5270 of our spectrum of NGC 4478 but this is unlikely to be the cause of the discrepancy here since no evidence of unsuccessful bad pixel correction was found. Since Fe5270, Fe5335, and $\langle Fe \rangle$ are critical indices in the study of stellar populations, it has to be noted that our results are based on our data. At the galaxy centre there is a small difference between our value of the Mg_2 index and that of Gorgas et al. (1990), while agreement within the errors is found for data at all other radii. The central measurements of the line-strength indices of González (1993) and Trager et al. (1998) are in good agreement with our measurements.




Figure 6. Comparison of our line-strength index data for NGC 4478 measured as a function of radius of the major axis (shown by filled triangles symbols), the data Gorgas et al. (1990, *open squares*) and Peletier (1989, *asterisks*). The central measurements of González (1993, *pentagon*) and Trager et al. (1998, *hexagon*) are shown for representative radii. Error bars are provided for data at the innermost and outermost radii and the central measurements.

4 AGES, METALLICITIES AND α/Fe OVERABUNDANCES

In the following we indicate the combined Magnesium-Iron index with

$$[MgFe]' = \sqrt{Mg\ b(0.72 \times Fe5270 + 0.28 \times Fe5335)} \quad (5)$$

newly defined by TMB. This index is completely independent of the $[\alpha/Fe]$ overabundance and hence serves best as a metallicity tracer.

We show in the top panels of Figure 7 the line strengths of the $H\beta$ and $[MgFe]'$ indices at different positions in the galaxies NGC 4458 and NGC 4478 with the grid of models of TMB. No clear trend is observed for NGC 4458. NGC 4478 shows a negative gradient in both the indices $H\beta$ and $[MgFe]'$. We show in bottom panels of Figure 7 the line strength of the $Mg\ b$ and $\langle Fe \rangle$ indices at different positions in the galaxies NGC 4458 and NGC 4478. For both galaxies, both indices $\langle Fe \rangle$ and $Mg\ b$ decrease outward on the mean.

Using stellar population models with variable element abundance ratios from TMB we derive average ages, metallicities and $[\alpha/Fe]$ ratios as a function of the position in the galaxies, following the procedure of Mehlert et al. (2003). Figure 8 translates the trends observed in Figure 7 as follows. NGC 4458 is uniformly old ($t \approx 15$ Gyr), has overall low metal content ($[Z/H] \approx -0.2$) and overabundance ($[\alpha/Fe] \approx 0.3$). For NGC 4478 instead the central regions ($R < 0.1R_e$) are younger ($t \approx 6$ Gyr), more metal rich ($[Z/H] \approx 0.35$) and less overabundant ($[\alpha/Fe] \approx 0.2$) than the outer regions, where old ages ($t \approx 15$ Gyr), low metallicities ($[Z/H] \approx 0.1$) and high overabundances ($[\alpha/Fe] \approx 0.3$) are observed on both the minor and the major axis. The size of the error bars shows that the decreases in age and overabundance in the cen-




Figure 5. The radial profiles of the line-strength indices measured along the major and minor axis of NGC 4458 (*left panels*) and NGC 4478 (*right panels*). Squares refer to the value of line-strength indices measured at the galaxy centre.

tral region are significant. Finally, note that within the errors most of the ages shown in Fig. 8 are compatible with the age of Universe favoured by WMAP (Spergel et al. 2003). In the following we will interpret the radial variations described above as the result of the superposition of the central discs detected photometrically in Section 2.2 and the main body of the galaxy. Moreover, we will assume that the main body of the galaxy does not have a $[\alpha/\text{Fe}]$ gradient, in agreement with the results of Mehlert et al. (2003). NGC 4458 has properties similar to giant ellipticals with decoupled cores. Prototypes of decoupled cores in giant ellipticals are as old as and metal richer than the rest of the galaxy, with high and approximately constant overabundance, as in the case of NGC 4816 and IC 4051 (Mehlert et al. 1998), or NGC 4365 (Surma & Bender 1995; Davies et al. 2001). In NGC 4458 the inner disc and the main body of the galaxy appear to have the same stellar populations, with a metallicity gradient for the main body of the galaxy slightly weaker than the cases discussed above. By contrast, the case of NGC 4478 is intriguing. The inner, cold disc is younger, richer in metals and less overabundant than the main body of the galaxy. We tested this conclusion with a composite stellar population model, summing 10% of a young ($t = 1.65$ Gyr) and metal rich ($Z = 3Z_{\odot}$) component with solar α/Fe ratios, and 90% of an old ($t = 12$ Gyr) metal rich ($Z = 2Z_{\odot}$) overabundant ($[\alpha/\text{Fe}] = 0.4$) population, and recovering the observed central values of the line indices. Given the errors of the available spectroscopic data and their spatial resolution, we refrained from a detailed study of the allowed range of alternative solutions, that only spectroscopy at HST resolution can pin down.

Of course, higher than solar α/Fe ratios for the disk are allowed if we relax the assumption that the α/Fe ratio of the main body of NGC 4478 is constant. Using these age and metallicity estimates and the models of Maraston (1998), we compute that the mass-to-light ratios of the inner discs are $M/L_V = 7.2 M_{\odot}/L_{\odot}$ for NGC 4458 (with a range from 5 to 10 judged from the range of ages and metallicities shown in Fig. 8) and $M/L_V = 1.3 M_{\odot}/L_{\odot}$ for NGC 4478. In this case, given the limitations of the spectroscopic data discussed above, the uncertainties are probably as large

as a factor 2. Combined with the luminosities given in Table 1, this gives masses of $M = 3.4 \times 10^7 M_{\odot}$ and $M = 2.5 \times 10^7 M_{\odot}$, respectively. The kinematic and photometric properties discussed above argue for the presence of (cold) discs at the centres of NGC 4458 and NGC 4478, and therefore for a formation scenario through the dissipational collapse of a gas rich object. The observed kinematic signature of counter-rotation in NGC 4458 argues in addition for an external origin with decoupled angular momentum. The absence of gradients in the stellar population of NGC 4458 indicates that the formation of the inner cold structure happened at the same time as the main body of the galaxy. The younger age and low overabundance of the central structure of NGC 4478 is indicative of a prolonged star formation history (Thomas, Greggio & Bender 1999), typical of an undisturbed disc-like, gas-rich (possibly pre-enriched) structure. This is consistent with the absence of counter-rotation.

5 CONCLUSIONS

We have investigated the properties of the central regions of the low-luminosity ellipticals NGC 4458 and NGC 4478, finding that:

(i) HST archive images of the two galaxies show the presence of central discs. They have scale lengths and face-on central surface brightnesses which fit on μ_0^c-h relation for galaxy discs. For NGC 4458 these parameters are typical of NSDs, while for NGC 4478 they are intermediate between the values for NSDs and the discs of discy ellipticals.

(ii) Measurements presented by Halliday et al. (2001) along the major and minor axis of the galaxies revealed the presence of decoupled kinematics on the radial scales of the NSDs. A counter-rotating central structure is present in NGC 4458, a cold rotating component in NGC 4478.

(iii) The Lick/IDS absorption line-strength indices $H\beta$, $Mg\ b$ and $\langle\text{Fe}\rangle$ measured as a function of radius along the major and the minor axis of the galaxies allow us to estimate the age, metallicity




Figure 7. *Top panels:* The line strength of the H β and [MgFe]' indices at different positions in the galaxies NGC 4458 (*left panel*) and NGC 4478 (*right panel*). The grid shows the models of Thomas et al. (2003) for different metallicities [Z/H] and ages. *Bottom panels:* The line strength of the Mg b and ⟨Fe⟩ indices at different positions in the galaxies NGC 4458 (*left panel*) and NGC 4478 (*right panel*). In all panels the *circles* show the major axis, the *triangles* the minor axis, the *filled symbols* the galaxy centres. The grid shows the models of Thomas et al. (2003) for different overabundances [α /Fe] and metallicities [Z/H], with age 12 Gyr.

and overabundance of the stellar populations of the two galaxies using simple stellar population models.

(iv) The radial variations of the derived quantities constrain the stellar population properties of the central discs. The NSD of NGC 4458 has an estimated stellar mass of $3.4 \times 10^7 M_{\odot}$ and is similar to the decoupled cores of bright ellipticals, being as old as and richer in metals than the rest of the galaxy, with high and approximately constant overabundance. The cold central disc of NGC 4478 has a similar estimated stellar mass of $2.5 \times 10^7 M_{\odot}$, but is younger, richer in metals and less overabundant than the main body of the galaxy.

(v) The nearly solar α -element abundance of the central disc of NGC 4478 indicates a prolonged star formation history, typical of an undisturbed disc-like, gas-rich (possibly pre-enriched) structure.

ACKNOWLEDGMENTS

Based on observations made with the NASA/ESA Hubble Space Telescope, obtained from the data archive at the Space Telescope

Institute. STScI is operated by the association of Universities for Research in Astronomy, Inc. under the NASA contract NAS 5-26555. We thank the observing committee of the CfA for their generous award of telescope time to obtain the spectroscopic observations presented here. The spectroscopic absorption line-strength measurements presented were derived as part of the Ph.D. thesis research of CH. CH gratefully acknowledges the support of a PPARC studentship at the University of Durham and the grant support of the Italian National Research Centre (grant CNRG00887) and a grant from the Fondo per gli Investimenti della Ricerca di Base of the Italian Ministry of Education, University and Research (grant RBAU018Y7E). We acknowledge A. W. Graham for useful discussion. LM gratefully acknowledges the support of an ESO studentship at the ESO Research Facilities in Santiago. RPS thanks the financial support by the DFG grant SFB375. This research has made use of the Lyon-Meudon Extragalactic Database (LED), NASA/IPAC Extragalactic Database (NED) and Starlink facilities.

REFERENCES

Arnold, R., de Zeeuw, P. T., Hunter, C., 1994, MNRAS, 271, 984A




Figure 8. The age (*top panels*), metallicity $[Z/H]$ (*middle panels*), and overabundance $[\alpha/Fe]$ profiles (*bottom panels*) as a function of the distance from the centre in units of half-luminosity radius derived from the $H\beta$, (Fe) and Mg b values using the models of Thomas et al. (2003). The *filled symbols* refer to the major axis of the galaxies, the *open symbols* to the minor axis.

Bender, R., Surma, P., Doebereiner, S., Moellenhoff, C., & Madejsky, R., 1989, *A&A*, 217, 35
 Bertola, F. & Corsini, E. M. 1999, *IAU Symp.* 186: Galaxy Interactions at Low and High Redshift, 186, 149
 Bertola, F., Corsini, E. M., Vega Beltrán, J. C., Pizzella, A., Sarzi, M., Cappellari, M., & Funes, J. G., 1999, *ApJL*, 519, L127
 Balcells, M., Graham, A. W., Domínguez-Palmero, L., & Peletier, R. F., 2003, *ApJL*, 582, L79
 Balcells, M., Graham, A. W., & Peletier, R. F., 2004, *ApJ*, submitted (astro-ph/0404381)
 Böker, T., Laine, S., van der Marel, R. P., Sarzi, M., Rix, H., Ho, L. C., & Shields, J. C. 2002, *AJ*, 123, 1389
 Bruzual, G. & Charlot, S., 2003, *MNRAS*, 344, 1000
 Cappellari, M. 2002, *MNRAS*, 333, 400
 Caon, N., Capaccioli, M., & Rampazzo, R., 1990, *A&AS*, 86, 429
 Carollo, C. M., Danziger, I. J., Rich, R. M., & Chen, X., 1997, *ApJ*, 491, 545
 Davies, R. L., Sadler, E. M., and Peletier, R. F., 1993, *MNRAS*, 262, 650
 Davies, R.L., Kuntschner, H., Emsellem, E., Bacon, R., Bureau, M., Carollo, C.M., Copin, Y., Miller, B.W., Monnet, G., Peletier, R.F., Verolme, E.K., de Zeeuw, P.T., 2001, *ApJL*, 548, L33
 de Vaucouleurs, G., de Vaucouleurs, A., Corwin, J. R., Buta, R. J., Paturel, G., & Fouque, P. 1991, *Third Reference Catalogue of Bright Galaxies (RC3)*
 Faber, S. M., Friel, E. D., Burstein, D., and Gaskell, C. M., 1985,

ApJS, 57, 711

Faber, S. M., Tremaine, Scott, Ajhar, Edward A., Byun, Yong-Ik, Dressler, Alan, Gebhardt, Karl, Grillmair, Carl, Kormendy, John, Lauer, Tod R., Richstone, Douglas, 1997, *AJ*, 114, 1771
 Ferrarese, L., van den Bosch, F. C., Ford, H. C., Jaffe, W., O'Connell, R. W., 1994, *AJ*, 108, 1598F
 González, J. J. 1993, *Line Strength Gradients and Kinematic Profiles in Elliptical Galaxies*, PhD thesis, University of California
 Gorgas, J., Efstathiou, G., & Aragón-Salamanca, A. A., 1990, *MNRAS*, 245, 217
 Graham, A. W. 2001, *MNRAS*, 326, 543
 Graham, A. W. & Guzmán, R. 2003, *AJ*, 125, 2936
 Halliday, C. 1999, *Low-Luminosity Elliptical Galaxies*, PhD thesis, University of Durham
 Halliday, C., Davies, R. L., Kuntschner, H., Bender, R., Birkinshaw, M., Saglia, R. P., & Baggley, G. 2001, *MNRAS*, 326, 473
 Haynes, M.P., Giovanelli, R., Chamaraux, P., da Costa, L.N., Freudling, W., Salzer, J.J., Wegner, G., 1997, *AJ*, 117, 2039
 Holley-Bockelmann, K. & Richstone, D. O., 2000, *ApJ*, 531, 232
 Holtzman, J. A. et al. 1995a, *PASP*, 107, 156
 Holtzman, J. A., Burrows, C. J., Casertano, S., Hester, J. J., Trauger, J. T., Watson, A. M., & Worthey, G. 1995b, *PASP*, 107, 1065
 Jaffe, W., Ford, H. C., O'Connell, R. W., van den Bosch, F. C., Ferrarese, L., 1994, *AJ*, 108, 1567J
 Jedrzejewski, R., 1987, *MNRAS*, 226, 747J
 Kinney, A. L., Calzetti, D., Bohlin, R. C., McQuade, K., Storchi-Bergmann, T., & Schmitt, H. R. 1996, *ApJ*, 467, 38
 Kobayashi, C. & Arimoto, N., 1999, *ApJ*, 527, 573
 Kormendy, J., 1984, *ApJ*, 287, 577
 Kormendy, J., Byun, Y.; Ajhar, E. A.; Lauer, T. R.; Dressler, A.; Faber, S. M.; Grillmair, C.; Gebhardt, K.; Richstone, D.; Tremaine, S., 1996, *IAU Symp.* 171: New Light on Galaxy Evolution, 171, 105
 Kormendy, J. & Bender, R., 1996, *ApJL*, 464, L119
 Kormendy, J. & Gebhardt, K., 2001, *AIP Conf. Proc.* 586: 20th Texas Symposium on Relativistic Astrophysics, 586, 363
 Krist, J., Hook, R. 1999, *STIS Instrument Handbook Version 4.0* (Baltimore: STScI)
 Lauer, T. R., Ajhar, E. A., Byun, Y.-I., Dressler, A., Faber, S. M., Grillmair, C., Kormendy, J., Richstone, D., Tremaine, S., 1995, *AJ*, 110, 2622
 Maraston, C., 1998, *MNRAS*, 300, 872
 Mehlert, D., Saglia, R.P., Bender, R., Wegner, G., 1998, *A&A*, 332, 33
 Mehlert, D., Saglia, R. P., Bender, R., & Wegner, G., 2000, *A&AS*, 141, 449
 Mehlert, D., Thomas, D., Saglia, R.P., Bender, R., Wegner, G., 2003, *A&A*, 407, 423
 Michard, R., 1985, *A&AS*, 59, 205
 Monnet, G., Bacon, R. & Emsellem, E. 1992, *A&A*, 253, 366
 Peletier, R. F. 1989, PhD thesis, University of Groningen
 Peletier, R. F., Davies, R. L., Illingworth, G. D., Davis, L. E., & Cawson, M., 1990, *AJ*, 100, 1091
 Pizzella, A., Corsini, E. M., Morelli, L., Sarzi, M., Scarlata, C., Stiavelli, M., & Bertola, F. 2002, *ApJ*, 573, 131
 Press, W. H., Teukolsky, S. A., Vetterling, W. T., & Flannery, B. P. 1992, *Numerical recipes in FORTRAN. The art of scientific computing* (Cambridge: University Press)
 Prugniel, P., Nieto, J.-L., & Simien, F., 1987, *A&A*, 173, 49
 Ravindranath, S., Ho, L. C., Peng, C. Y., Filippenko, A. V., Sargent, W. L. W., 2001, *AJ*, 122, 653

- Rest, A., van den Bosch, F. C., Jaffe, W., Tran, H., Tsvetanov, Z., Ford, H. C., Davies, J., & Schafer, J., 2001, *AJ*, 121, 2431
- Rix, H. W. and White, S. D. M., 1992, *MNRAS*, 254, 389
- Sandage, A. & Tammann, G. A. 1981, Revised Shapley-Ames Catalog of Bright Galaxies, (Washington: Carnegie Institution) (RSA)
- Scorza, C. & Bender, R. 1995, *A&A*, 293, 20 (SB95)
- Scorza, C. & van den Bosch, F. C. 1998, *MNRAS*, 300, 469
- Seifert, W. & Scorza, C. 1996, *A&A*, 310, 75 (SS96)
- Spergel, D. N., Verde, L., Peiris, H. V., Komatsu, E., Nolta, M. R., Bennett, C. L., Halpern, M., Hinshaw, G., Jarosik, N., Kogut, A., Limon, M., Meyer, S. S., Page, L., Tucker, G. S., Weiland, J. L., Wollack, E., & Wright, E. L. 2003, *ApJS*, 148, 175
- Surma, P. & Bender, R. 1995, *A&A*, 298, 405
- Tantalo, R., Chiosi, C., & Bressan, A., 1998, *A&A*, 333, 419
- Thomas, D., Greggio, L, Bender, R., 1999 *MNRAS*, 302, 537
- Thomas D., Maraston, C., Bender R. 2003, *MNRAS*, 339, 897 (TMB)
- Tomita, A., Aoki, K., Watanabe, M., Takata, T., Ichikawa, S., 2000, *AJ*, 120, 123
- Trager, S. C. 1997 The Stellar Population Histories of Elliptical Galaxies, PhD thesis, University of California
- Trager, S. C., Worthey, G., Faber, S. M., Burstein, D., & González, J. J. 1998, *ApJS*, 116, 1
- Tran, H. D., Tsvetanov, Z., Ford, H. C., Davies, J., Jaffe, W., van den Bosch, F. C., Rest, A., 2001, *AJ*, 121, 2928
- Trujillo, I., Erwin, P., Ramos, A. A., & Graham, A. W. 2004, *AJ*, 127, 1917
- Tully, R. B. 1988, *Nearby Galaxy Catalog*, (Cambridge: Cambridge University Press)
- van den Bosch, F. C., Ferrarese, L., Jaffe, W. Ford, H.C., O'Connell, R.W., 1994, *AJ*, 108, 1579
- van den Bosch, F. C. 1998, *ApJ*, 507, 601
- van den Bosch, F. C. & Emsellem, E., 1998, *MNRAS*, 298, 267
- van den Bosch, F. C., Jaffe, W. & van der Marel, R. P. 1998, *MNRAS*, 293, 343
- van Dokkum, P. G. & Franx, M., 1995, *AJ*, 110, 2027V
- Vazdekis, A., 1999, *ApJ*, 513, 224
- Worthey, G., 1994, *ApJS*, 95,107